Constraints on the Primordial Magnetic Field from σ_8

Dai G. Yamazaki[1],* Kiyotomo Ichiki[2], Toshitaka Kajino[1,3], and Grant. J. Mathews[4]
[1]*National Astronomical Observatory of Japan 2-21-1 Osawa, Mitaka, Tokyo 181-8588, Japan*
[2]*Research Center for the Early Universe, School of Science, the University of Tokyo, Hongo 7-3-1, Bunkyo-ku, Tokyo 113-0033, Japan*
[3] *Department of Astronomy, School of Science, The University of Tokyo, Hongo 7-3-1, Bunkyo-ku, Tokyo 113-0033, Japan and*
[4]*Center for Astrophysics, Department of Physics, University of Notre Dame, Notre Dame, IN 46556, U.S.A.*
(Dated: November 13, 2008)

A primordial magnetic field (PMF) can affect the evolution of density field fluctuations in the early universe. In this paper we constrain the PMF amplitude B_λ and power spectral index $n_{\rm B}$ by comparing calculated density field fluctuations with observational data, i.e. the number density fluctuation of galaxies. We show that the observational constraints on cosmological density fluctuations, as parameterized by σ_8, lead to strong constraints on the amplitude and spectral index of the PMF.

PACS numbers: 98.62.En,98.70.Vc

arXiv:0811.2221v1 [astro-ph] 13 Nov 2008

I. INTRODUCTION

Magnetic fields have been observed [1, 2, 3, 4] in clusters of galaxies with a strength of $0.1 - 1.0$ μ G. One possible explanation for such magnetic fields in galactic clusters is the existence of a primordial magnetic field (PMF) of order 1 nG whose field lines collapse as structure forms. Therefore, recently, the origin and amplification mechanism of the PMF in the scale of galaxy cluster have been proposed and studied intensively by a number of authors [5, 6, 7, 8, 9, 10, 11, 12, 13, 14]. The origin and detection of the PMF is, hence, a subject of considerable interest in modern cosmology. Moreover, the PMF could influence a variety of phenomena in the early universe [15, 16, 17, 18] such as the cosmic microwave background (CMB) [10, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36], and the formation of large-scale structure (LSS) [37, 38, 39, 40, 41].

If dynamically significant large-scale magnetic fields were present in the early universe, they would have affected the formation and evolution of the observed structure. Thus, some signatures of the existence of a PMF should be apparent in the presently observed cosmic structure.

In this regard, the alternative normalization parameter σ_8 is of particular interest. It is defined [42] as the root-mean-square of the matter density fluctuations in a comoving sphere of radius $8h^{-1}$ Mpc. It is determined by a weighted integral of the matter power spectrum. Observations which determine σ_8 provide information about the physical processes affecting the evolution of density-field fluctuations and the formation of structure on the cosmological scales. The mechanisms by which a PMF can affect the density field fluctuations on cosmological scales has been described in our previous work [37]. Of course, σ_8 is also affected by the presence of a PMF. In this article we show that by considering the effect of a PMF on σ_8 and comparing theoretically estimated values for σ_8 with the observed range, we can obtain not only insight into the underlying physical processes of density field fluctuations in the presence of a PMF, but also place constraints on the amplitude and spectral index of the PMF.

II. THE MODEL

We use the isocurvature magnetized initial conditions with adiabatic relations for the fluids evolution of primary density perturbations and in the presence of a PMF. For the present purposes we fix the cosmological parameters to those of the best-fit flat ΛCDM model as given in Ref. [43], i.e. $h = 0.719$, $\Omega_b h^2 = 0.02273$, $\Omega_c h^2 = 0.1099$, $n_S = 0.963$, and $\tau_c = 0.087$, where h denotes the Hubble parameter in units of 100 km s^{-1}Mpc^{-1}, Ω_b and Ω_c are the baryon and cold dark matter densities in units of the critical density, n_S is the spectral index of the primordial scalar fluctuations, and τ_c is the optical depth for Compton scattering. We use natural units $c = \hbar = 1$.

*URL: `http://th.nao.ac.jp/~yamazaki/`; Electronic address: yamazaki@th.nao.ac.jp

A. Primordial Magnetic Field

Before recombination, Thomson scattering between photons and electrons along with Coulomb interactions between electrons and baryons were sufficiently rapid to ensure that the photon-baryon plasma behaved as a single tightly coupled fluid. Since the trajectories of plasma particles are bent by Lorentz forces in a magnetic field, photons are indirectly influenced by the magnetic field through Thomson scattering. The energy density of the magnetic field can be treated as a first order perturbation upon a flat Friedmann-Robertson-Walker (FRW) background metric. In this linear approximation[44], the magnetic field evolves as a stiff source. Therefore, we can discard all back reaction terms from the magnetohydrodynamic (MHD) fluid onto the field itself.

B. Power Spectrum from the PMF

During the epochs of interest here, the conductivity of the primordial plasma is very large, and the PMF is "frozen-in" to a very good approximation[25]. Furthermore, we can neglect the electric field, i.e. $E \sim 0$, and can decouple the time evolution of the magnetic field from its spatial dependence, i.e. $\mathbf{B}(\mathbf{x},\tau) = \mathbf{B}(\mathbf{x})/a^2$ for very large scales. This leads to the following simplified electromagnetic energy-momentum tensor,

$$T^{00}{}_{[\mathrm{EM}]}(\mathbf{x},\tau) = \frac{B(\mathbf{x})^2}{8\pi a^6} \quad , \tag{1}$$

$$T^{i0}{}_{[\mathrm{EM}]}(\mathbf{x},\tau) = T^{0k}{}_{[\mathrm{EM}]}(\mathbf{x},\tau) = 0 \quad , \tag{2}$$

$$-T^{ik}{}_{[\mathrm{EM}]}(\mathbf{x},\tau) = \sigma_{\mathrm{B}}^{ik} = \frac{1}{8\pi a^6}\left\{2B^i(\mathbf{x})B^k(\mathbf{x}) - \delta^{ik}B(\mathbf{x})^2\right\} \quad . \tag{3}$$

We assume that the initial PMF is statistically homogeneous, isotropic and random. For such a magnetic field, the power spectrum can be taken as a power-law $P(k) \propto k^{n_B}$ [25, 31] where n_B is the spectral index which can be either negative or positive depending upon the physical processes of the field creation. From ref. [25], a two-point correlation function for the PMF can be defined by

$$\left\langle B^i(\boldsymbol{k})B^{j*}(\boldsymbol{k}')\right\rangle \;=\; \frac{(2\pi)^{n_B+8}}{2k_\lambda^{n+3}}\frac{B_\lambda^2}{\Gamma\left(\frac{n_B+3}{2}\right)}k^{n_B}P^{ij}(k)\delta(\boldsymbol{k}-\boldsymbol{k}'), \quad k < k_C, \tag{4}$$

where

$$P^{ij}(k) \;=\; \delta^{ij} - \frac{k^i k^j}{k^2} \quad . \tag{5}$$

Here, B_λ is the comoving mean magnetic-field amplitude obtained by smoothing over a Gaussian sphere of comoving radius λ, and $k_\lambda \equiv 2\pi/\lambda$ (with $\lambda = 1$ Mpc in this paper). Hereafter, we work in k-space and denote all quantities by their Fourier transform convention

$$F(\boldsymbol{k}) = \int d^3x \exp(i\boldsymbol{k}\cdot\boldsymbol{x})F(\boldsymbol{x}). \tag{6}$$

The cutoff wave number k_C in the magnetic power spectrum is defined by [45, 46],

$$k_C^{-5-n_B}(\tau) = \begin{cases} \frac{B_\lambda^2 k_\lambda^{-n_B-3}}{4\pi(\rho+p)}\int_0^\tau d\tau' \frac{l_\gamma}{a}, & \tau < \tau_{\mathrm{dec}} \\ k_C^{-5-n_B}(\tau_{\mathrm{dec}}), & \tau > \tau_{\mathrm{dec}} \end{cases} \quad , \tag{7}$$

where l_γ is the photon mean free path, and τ_{dec} is the conformal time at the epoch of photon-baryon decoupling.

We obtain power spectra for the PMF energy density and the Lorentz force for the scalar mode, respectively, as follows

$$|E_{[\mathrm{EM:S}]}(\boldsymbol{k},\tau)|^2\delta(\boldsymbol{k}-\boldsymbol{k}') = \frac{1}{(2\pi)^3}\left\langle T(\boldsymbol{k},\tau)_{[\mathrm{EM:S1}]}T^*(\boldsymbol{k}',\tau)_{[\mathrm{EM:S1}]}\right\rangle \quad , \tag{8}$$

and

$$\begin{aligned} |\Pi_{[\mathrm{EM:S}]}(\boldsymbol{k},\tau)|^2\delta(\boldsymbol{k}-\boldsymbol{k}') = \quad & \frac{1}{(2\pi)^3}\left\langle \left(T(\boldsymbol{k},\tau)_{[\mathrm{EM:S1}]} - T(\boldsymbol{k},\tau)_{[\mathrm{EM:S2}]}\right)\right. \\ & \left.\times\left(T^*(\boldsymbol{k}',\tau)_{[\mathrm{EM:S1}]} - T^*(\boldsymbol{k}',\tau)_{[\mathrm{EM:S2}]}\right)\right\rangle \ , \end{aligned} \tag{9}$$

where S1 and S2 in the subscripts of the energy-momentum tensor denote the PMF energy density and pressure. An explicit expression can be obtained for the ensemble averages which are used to evaluate the above spectra. In the case of a power law stochastic magnetic field we have[31, 37, 47, 48],

$$\langle T(\boldsymbol{k},\tau)_{[\mathrm{EM:S1}]}T^{*}(\boldsymbol{p},\tau)_{[\mathrm{EM:S1}]}\rangle = \frac{1}{2^4(2\pi)^8a^8}\left\{\frac{(2\pi)^{n+8}}{2k_\lambda^{n+3}}\frac{B_\lambda^2}{\Gamma\left(\frac{n+3}{2}\right)}\right\}^2 \times\int d^3k'k'^n|\boldsymbol{k}-\boldsymbol{k}'|^n\left\{1+\frac{\{\boldsymbol{k}'\cdot(\boldsymbol{k}-\boldsymbol{k}')\}^2}{k'^2|\boldsymbol{k}-\boldsymbol{k}'|^2}\right\}\delta(\boldsymbol{k}-\boldsymbol{p})\ . \tag{10}$$

The two-point correlation function for the Lorentz force is given by

$$\langle T(\boldsymbol{k},\tau)_{[\mathrm{EM:S1}]}T(\boldsymbol{p},\tau)^{*}_{[\mathrm{EM:S1}]}\rangle = \frac{1}{2^3(2\pi)^7a^8}\left\{\frac{(2\pi)^{n+8}}{2k_\lambda^{n+3}}\frac{B_\lambda^2}{\Gamma\left(\frac{n+3}{2}\right)}\right\}^2 \times\int dk'k'^{n+2}\int_{-1}^{1}d\mathcal{C}|\boldsymbol{k}-\boldsymbol{k}'|^{n-2}\left\{(1+\mathcal{C}^2)k^2-4kk'\mathcal{C}+2k'^2\right\}\delta(\boldsymbol{k}-\boldsymbol{p})\ \ . \tag{11}$$

Here, we define $\mathcal{C}$ as

$$\mathcal{C}=\cos c=\hat{\boldsymbol{k}}\cdot\hat{\boldsymbol{k}}'=\frac{\boldsymbol{k}'\cdot\boldsymbol{k}}{k'k}\ . \tag{12}$$

Almost all previous works have set the terms which include $\mathcal{C}$ in the middle parenthesis to unity. In this paper, however, we evaluate Eq. (11) explicitly using integration by parts. In this way we obtain the following equation.

$$\begin{aligned}\langle T(\boldsymbol{k},\tau)_{[\mathrm{EM:S1}]}T^{*}(\boldsymbol{k},\tau)_{[\mathrm{EM:S1}]}\rangle &= \frac{1}{8\pi a^8}\left\{\frac{(2\pi)^{n_B+5}}{2k_\lambda^{n_B+3}}\frac{B_\lambda^2}{\Gamma\left(\frac{n_B+3}{2}\right)}\right\}^2\\ &\times\int dk'k'^{n_B+2}\left[\frac{n_B^2+4n_B+1}{kk'n_B(n_B+2)(n_B+4)}\left\{(k+k')^{n_B+2}-|k-k'|^{n_B+2}\right\}\right.\\ &-\frac{1}{k'^2n_B(n_B+4)}\left\{|k-k'|^{n_B+2}+|k+k'|^{n_B+2}\right\}\\ &\left.+\frac{k}{k'^3n_B(n_B+2)(n_B+4)}\left\{(k+k')^{n_B+2}-|k-k'|^{n_B+2}\right\}\right].\end{aligned} \tag{13}$$

A similar derivation leads to the power spectrum of the PMF tension as well as the power spectrum of the correlation between pressure and tension as follows,

$$\begin{aligned}\left\langle T_{[\mathrm{EM:S2}]}(\boldsymbol{k})T^{*}_{[\mathrm{EM:S2}]}(\boldsymbol{k})\right\rangle &= \frac{1}{2\pi a^8}\left\{\frac{(2\pi)^{n_B+5}}{2k_\lambda^{n_B+3}}\frac{B_\lambda^2}{\Gamma\left(\frac{n_B+3}{2}\right)}\right\}^2\\ &\times\int dk'k'^{n_B+4}\frac{4}{(kk')^3n_B(n_B+2)(n_B+4)}\left[\ \left\{(k+k')^{n_B+4}-|k-k'|^{n_B+4}\right\}\right.\\ &-\frac{3}{(kk')(n_B+6)}\left\{|k-k'|^{n_B+6}+(k+k')^{n_B+6}\right\}\\ &\left.+\frac{3}{(kk')^2(n_B+6)(n_B+8)}\left\{(k+k')^{n_B+8}-|k-k'|^{n_B+8}\right\}\right]\ ,\end{aligned} \tag{14}$$

and

$$
\begin{aligned}
&\langle T_{[\mathrm{EM:S1}]}(\boldsymbol{k})T^{*}_{[\mathrm{EM:S2}]}(\boldsymbol{k})\rangle + \langle T_{[\mathrm{EM:S2}]}(\boldsymbol{k})T^{*}_{[\mathrm{EM:S1}]}(\boldsymbol{k})\rangle \\
&= \frac{1}{2\pi a^8}\left\{\frac{(2\pi)^{n_B+5}}{2k_\lambda^{n_B+3}}\frac{B_\lambda^2}{\Gamma\left(\frac{n_B+3}{2}\right)}\right\}^2 \\
&\quad\times\int dk' k'^{n_B+3}\left[\frac{1}{(kk')^2 n_B(n_B+2)}\left\{(k+k')^{n_B+3}-|k-k'|^{n_B+3}\right\}\right. \\
&\quad-\frac{3}{k^2k'^3 n_B(n_B+2)(n_B+4)}\left\{|k-k'|^{n_B+4}+(k+k')^{n_B+4}\right\} \\
&\quad-\frac{1}{k^3k'^2 n_B(n_B+2)(n_B+4)}\left\{(k+k')^{n_B+4}-|k-k'|^{n_B+4}\right\} \\
&\quad\left.+\frac{3}{k^3k'^4 n_B(n_B+2)(n_B+4)(n_B+6)}\left\{(k+k')^{n_B+6}-|k-k'|^{n_B+6}\right\}\right] \quad .
\end{aligned}
\tag{15}
$$

For this article we have constructed a numerical program, "PriME: Program for primordial Magnetic Effects", with which we can evaluate the PMF source power spectrum using the numerical method described in Refs. [23, 37, 49]. Using this, we can quantitatively evaluate the time evolution of the cut off scale and thereby reliably calculate the effects of the PMF.

III. EVOLUTION EQUATIONS

We now summarize the essential evolution equations for each mode.
For the scalar mode we obtain the following equations in k-space [35, 50, 51, 52, 53, 54]:

$$
k^2\phi + 3H(\dot{\phi}+H\psi) = 4\pi G a^2\left\{E_{[\mathrm{EM:S}]}(\boldsymbol{k},\tau)-\delta\rho_{\mathrm{tot}}\right\} \tag{16}
$$

$$
\begin{aligned}
k^2(\phi-\psi) &= -12\pi G a^2\left\{Z_{[\mathrm{EM:S}]}(\boldsymbol{k},\tau)-(\rho_\nu+P_\nu)\sigma_\nu\right\} \\
&= -12\pi G a^2\left\{\frac{1}{3}E_{[\mathrm{EM:S}]}(\boldsymbol{k},\tau)+\Pi_{[\mathrm{EM:S}]}(\boldsymbol{k},\tau)-(\rho_\nu+P_\nu)\sigma_\nu\right\} \quad ,
\end{aligned}
\tag{17}
$$

$$
\begin{aligned}
\dot{\delta}^{(\mathrm{S})} &= -(1+w)\left(v^{(\mathrm{S})}+3\dot{\phi}\right)-3H\left(\frac{\delta p}{\delta\rho}-w\right)\delta^{(\mathrm{S})} \\
&\quad-\frac{3}{8\pi\rho}\left\{\dot{E}_{[\mathrm{EM:S}]}(\boldsymbol{k},\tau)+6HE_{[\mathrm{EM:S}]}(\boldsymbol{k},\tau)\right\} \quad ,
\end{aligned}
\tag{18}
$$

$$
\begin{aligned}
\dot{v}^{(\mathrm{S})} &= -H(1-3w)v^{(\mathrm{S})}-\frac{\dot{w}}{1+w}v^{(\mathrm{S})}+\frac{\delta p}{\delta\rho}\frac{k^2\delta^{(\mathrm{S})}}{1+w}-k^2\sigma+k^2\psi \\
&\quad+k^2\frac{\Pi_{[\mathrm{EM:S}]}(\boldsymbol{k},\tau)}{4\pi\rho} \quad ,
\end{aligned}
\tag{19}
$$

where $Z_{[\mathrm{EM:S}]}\equiv\sigma_{\mathrm{B}}(\rho_\gamma+P_\gamma)$ and $w\equiv p/\rho$ is the usual equation of state parameter. Note that for the photon $\delta_\gamma^{(\mathrm{S})}=4\Theta_0^{(\mathrm{S})}$, and $v_\gamma^{(\mathrm{S})}=k\Theta_1^{(\mathrm{S})}$. Massless neutrinos obey Eqs. (18) and (19) as written without the Thomson coupling term. In the continuity and Euler relations (Eqs. 18 and 19) for the scalar mode, we can just add the energy density and pressure of the PMF to the energy density and pressure of the cosmic fluids. Since the baryon fluid behaves like a nonrelativistic fluid during the epoch of interest, we can neglect w and $\delta P_b^{(\mathrm{S})}/\delta\rho_b^{(\mathrm{S})}$, except for the acoustic term $c_s k^2\delta_b^{(\mathrm{S})}$. Also, the shear stress of the baryons is negligible [51]. Since we concentrate on scalar type perturbations in this paper, we do not consider the magneto-rotational instability from the shear stress of the PMF and baryon fluid [55].

By considering the Compton interaction between baryons and photons in equations (18) and (19) we obtain the

same form for the evolution equations of photons and baryons as in previous work [50, 51, 52, 53, 54].

$$
\begin{aligned}
k^2\phi + 3H(\dot{\phi} + H\psi) &= 4\pi G a^2 \left\{ E_{[\mathrm{EM:S}]}(\boldsymbol{k},\tau) - \delta\rho_{\mathrm{tot}} \right\} && (20)\\
k^2(\phi - \psi) &= -12\pi G a^2 \left\{ Z_{[\mathrm{EM:S}]}(\boldsymbol{k},\tau) - (\rho_\nu + P_\nu)\sigma_\nu \right\} \\
&= -12\pi G a^2 \left\{ \frac{1}{3} E_{[\mathrm{EM:S}]}(\boldsymbol{k},\tau) + \Pi_{[\mathrm{EM:S}]}(\boldsymbol{k},\tau) - (\rho_\nu + P_\nu)\sigma_\nu \right\} \ , && (21)\\
\dot{\delta}^{(\mathrm{S})}_{\mathrm{CDM}} &= -v^{(\mathrm{S})}_{\mathrm{CDM}} + 3\dot{\phi} \ , && (22)\\
\dot{v}^{(\mathrm{S})}_{\mathrm{CDM}} &= -\frac{\dot{a}}{a} v^{(\mathrm{S})}_{\mathrm{CDM}} + k^2\psi \ , && (23)\\
\dot{\delta}^{(\mathrm{S})}_{\gamma} &= -\frac{4}{3} v^{(\mathrm{S})}_{\gamma} + 4\dot{\phi} \ , && (24)\\
\dot{\delta}^{(\mathrm{S})}_{\nu} &= -\frac{4}{3} v^{(\mathrm{S})}_{\nu} + 4\dot{\phi} \ , && (25)\\
\dot{v}^{(\mathrm{S})}_{\gamma} &= k^2 \left(\frac{1}{4}\delta^{(\mathrm{S})}_{\gamma} - \sigma_\gamma \right) + a n_e \sigma_T (v^{(\mathrm{S})}_{\mathrm{b}} - v^{(\mathrm{S})}_{\gamma}) \ + k^2\psi, && (26)\\
\dot{v}^{(\mathrm{S})}_{\nu} &= k^2 \left(\frac{1}{4}\delta^{(\mathrm{S})}_{\nu} - \sigma_\nu \right) + k^2\psi, && (27)\\
\dot{\delta}^{(\mathrm{S})}_{\mathrm{b}} &= -v^{(\mathrm{S})}_{\mathrm{b}} + 3\dot{\phi} \ , && (28)\\
\dot{v}^{(\mathrm{S})}_{\mathrm{b}} &= -\frac{\dot{a}}{a} v^{(\mathrm{S})}_{\mathrm{b}} + c_s^2 k^2 \delta^{(\mathrm{S})}_{\mathrm{b}} + \frac{4\bar{\rho}_\gamma}{3\bar{\rho}_{\mathrm{b}}} a n_e \sigma_T (v^{(\mathrm{S})}_{\gamma} - v^{(\mathrm{S})}_{\mathrm{b}}) + k^2\psi \\
&\quad + \frac{3}{4} k^2 \frac{\Pi_{[\mathrm{EM:S}]}(\boldsymbol{k},\tau)}{R\rho_\gamma} \ , && (29)
\end{aligned}
$$

where $R \equiv (3/4)(\rho_b/\rho_\gamma)$ is the inertial density ratio of baryons to photons, n_e is the free electron density, σ_T is the Thomson scattering cross section, and σ_γ of the second term on the right hand side of equation (26) is the shear stress of the photons with the PMF. Since $n_{\mathrm{B}} \lesssim 0$ is favored by constraints from the gravitational wave background [56], and the PMF effects are not influenced by the time evolution of the cut off scale k_C for this range of n_{B}, we can approximately set $E_{[\mathrm{EM:S}]} \propto a^{-4}$ in the following analysis.

A. Initial Conditions

We need to specify the initial perturbations for solving the evolution equations presented in the previous section. We start the solution at early times when the k modes of interest are still outside the horizon, i.e. the dimensionless parameter $k\tau \ll 1$. We consider only the radiation-dominated epoch since the numerical integration for all of the k modes of interest will start within this era. Baryons and photons are tightly coupled at this early time and the expansion rate is $H = \tau^{-1}$. We derive initial conditions for all of the modes utilizing the method of Refs. [35, 50, 51, 52, 53, 54]. We can assume that all density fields are zero initially, since the PMF only affects the velocity field of ionized baryons, via the Lorentz force, and the density fields are not directly affected by the PMF. In the radiation dominated epoch, photons and neutrinos are important in the energy-momentum tensor. The evolution equations for the photons and neutrinos are

$$
\begin{aligned}
\dot{\delta}^{(\mathrm{S})}_{\gamma} &= -\frac{4}{3} v^{(\mathrm{S})}_{\gamma} + 4\dot{\phi} \ , && (30)\\
\dot{\delta}^{(\mathrm{S})}_{\nu} &= -\frac{4}{3} v^{(\mathrm{S})}_{\nu} + 4\dot{\phi} \ , && (31)\\
\dot{v}^{(\mathrm{S})}_{\gamma} &= k^2 \frac{1}{4} \delta^{(\mathrm{S})}_{\gamma} + k^2\psi, && (32)\\
\dot{v}^{(\mathrm{S})}_{\nu} &= k^2 \left(\frac{1}{4}\delta^{(\mathrm{S})}_{\nu} - \sigma^{(\mathrm{S})}_{\nu} + k^2\psi \right), && (33)\\
\sigma^{(\mathrm{S})}_{\nu} &= \frac{4}{15} v^{(\mathrm{S})}_{\gamma}. && (34)
\end{aligned}
$$

Here, we have omitted higher multipole moments $\ell > 1$ for photons and $\ell > 2$ for neutrinos. At lowest order in $k\tau$, the initial conditions for Eqs.(20-29) are

$$\delta_\gamma^{(\mathrm{S})} = \delta_\nu^{(\mathrm{S})} = \frac{4}{3}\delta_b^{(\mathrm{S})} = \frac{4}{3}\delta_{\mathrm{CDM}}^{(\mathrm{S})} = R_\gamma R_{\mathrm{B}} + 4R_\gamma \frac{4\sigma_{\mathrm{B}} + R_\nu R_{\mathrm{B}}}{4R_\nu + 15}, \tag{35}$$

$$v_\gamma^{(\mathrm{S})} = v_b^{(\mathrm{S})} = v_{\mathrm{CDM}}^{(\mathrm{S})} = -\frac{19}{4}\frac{4\sigma_{\mathrm{B}} + R_\nu R_{\mathrm{B}}}{4R_\nu + 15}k^2\tau, \tag{36}$$

$$v_\nu^{(\mathrm{S})} = -\frac{15}{4}\frac{R_\gamma}{R_\nu}\frac{4\sigma_{\mathrm{B}} + R_\nu R_{\mathrm{B}}}{4R_\nu + 15}k^2\tau, \tag{37}$$

$$\sigma_\nu^{(\mathrm{S})} = -\frac{R_\gamma}{R_\nu}\sigma_{\mathrm{B}} + \frac{R_\gamma}{2R_\nu}\frac{4\sigma_{\mathrm{B}} + R_\nu R_{\mathrm{B}}}{4R_\nu + 15}k^2\tau^2, \tag{38}$$

$$\psi = -2\phi = -2R_\gamma \frac{R_\gamma}{2R_\nu}\frac{4\sigma_{\mathrm{B}} + R_\nu a R_{\mathrm{B}}}{4R_\nu + 15}\ , \tag{39}$$

where

$$R_\gamma \equiv \frac{\rho_\gamma}{\rho_\gamma + \rho_\nu},$$
$$R_\nu \equiv \frac{\rho_\nu}{\rho_\gamma + \rho_\nu},$$
$$R_{\mathrm{B}} \equiv \frac{E_{[\mathrm{EM:S}]}}{\rho_\gamma}\ .$$

IV. MATTER POWER SPECTRUM

Possible origins of the PMF have been studied by many authors, however, there is no consensus yet as to the origin of the PMF. Thus, we cannot know how the PMF correlates with the primordial density fluctuations. However, almost all previous works investigated the effects of a PMF on density perturbations under the assumption that there is no correlation between the PMF and the primordial density fluctuations [57]. However, in order to study the PMF effects in a more general manner, we introduce a parameter "s" which characterizes the correlation between the PMF and the primordial density fluctuations[35, 37]. In the linear approximation, the power spectra of the baryon ($P_{\mathrm{b}}(k)$) and CDM ($P_{\mathrm{CDM}}(k)$) density fluctuations in the presence of a PMF are then written,

$$\begin{aligned} P_{\mathrm{b}}(k) &= \left\langle \delta_{[\mathrm{b:FL}]}(k)\delta^*_{[\mathrm{b:FL}]}(k)\right\rangle + \left\langle \delta_{[\mathrm{b:PMF}]}(k)\delta^*_{[\mathrm{b:PMF}]}(k)\right\rangle \\ &+ 2\left\langle \delta_{[\mathrm{b:FL}]}(k)\delta^*_{[\mathrm{b:PMF}]}(k)\right\rangle, \end{aligned} \tag{40}$$

$$\begin{aligned} P_{\mathrm{CDM}}(k) &= \left\langle \delta_{[\mathrm{CDM:FL}]}(k)\delta^*_{[\mathrm{CDM:FL}]}(k)\right\rangle + \left\langle \delta_{[\mathrm{CDM:PMF}]}(k)\delta^*_{[\mathrm{CDM:PMF}]}(k)\right\rangle \\ &+ 2\left\langle \delta_{[\mathrm{CDM:FL}]}(k)\delta^*_{[\mathrm{CDM:PMF}]}(k)\right\rangle, \end{aligned} \tag{41}$$

where we normalize the cross correlation terms with the parameter s,

$$\left\langle \delta_{[\mathrm{b:FL}]}(k)\delta^*_{[\mathrm{b:PMF}]}(k)\right\rangle \equiv s\sqrt{\left\langle \delta_{[\mathrm{b:FL}]}(k)\delta^*_{[\mathrm{b:FL}]}(k)\right\rangle \left\langle \delta_{[\mathrm{b:PMF}]}(k)\delta^*_{[\mathrm{b:PMF}]}(k)\right\rangle}, \tag{42}$$

$$\left\langle \delta_{[\mathrm{CDM:FL}]}(k)\delta^*_{[\mathrm{CDM:PMF}]}(k)\right\rangle \equiv s\sqrt{\left\langle \delta_{[\mathrm{CDM:FL}]}(k)\delta^*_{[\mathrm{CDM:FL}]}(k)\right\rangle \left\langle \delta_{[\mathrm{CDM:PMF}]}(k)\delta^*_{[\mathrm{CDM:PMF}]}(k)\right\rangle}\ . \tag{43}$$

Here, δ_α, $\alpha \in ([\mathrm{b:FL}], [\mathrm{CDM:FL}])$ designate the baryon and CDM density fluctuations without the PMF respectively. Similarly, δ_β, $\beta \in ([\mathrm{b:PMF}], [\mathrm{CDM:PMF}])$ denote the baryon and CDM density fluctuations with the PMF included. When $0 < s \leq 1$, $s = 0$, or $-1 \leq s < 0$ in eqs.(42) and (43), one has positive, vanishing, or negative correlations, respectively.

The square root of the power spectrum functions for the Lorentz force $\Pi_{[\mathrm{EM:S}]}(\boldsymbol{k}, \tau)$ in Eq.(29) does not specify the sign. In other words, there is no information as to whether the magnetic pressure or the tension is dominant, and whether the directions of forces from them are the same or different. However, such information should be taken into account.

The Lorentz force term in Eq.(29) can be divided into two terms, the magnetic pressure and the tension. Their amplitudes are given by Eqs. (13) and (14), respectively. By comparing those equations, one can decide which of them is dominant in the Lorentz force term. We find that the former dominates when $n < -1.5$, while the latter dominates for $n > -1.5$. However, one cannot determine the relative signs of those two terms unless one also specifies a model for the generation of the PMF [37]. Thus, we must decompose the factors into various combinations, i.e.

$$s = s_{[\mathrm{LF}]} \times s_{[\mathrm{DF}]}, \tag{44}$$

where

$$s_{[\mathrm{LF}]} = \begin{cases} -1, & n < -1.5 \quad \text{(I)}, \\ -1, & n > -1.5 \quad \text{(II)}, \\ 1, & n > -1.5 \quad \text{(III)}, \end{cases} \tag{45}$$

and

$$\begin{aligned} 0 \;<\; s_{[\mathrm{DF}]} \;&\leq\; 1 \quad \text{(i)}, \\ s_{[\mathrm{DF}]} \;&=\; 0 \quad \text{(ii)}, \\ -1 \;\leq\; s_{[\mathrm{DF}]} \;&<\; 0 \quad \text{(iii)}. \end{aligned}$$

In the different regimes, $s_{[\mathrm{LF}]}$ represents either: (I) the pressure dominated case; (II) the tension dominated case, where the magnetic field pressure and tension forces act in the same direction; or (III) the tension dominated case, where the magnetic field pressure and tension forces act in the opposite direction. On the other hand, $s_{[\mathrm{DF}]}$ represents either: (i) a positive correlation between the matter and PMF distributions; (ii) no correlation; or (iii) a negative correlation. Thus, if $s < 0$, the matter and PMF distributions could be correlated positively ($s_{[\mathrm{DF}]} > 0$) and the PMF pressure dominates in the Lorentz term (for $n < -1.5$). Another possibility is that the matter and PMF distributions negatively correlate ($s_{[\mathrm{DF}]} < 0$) and the PMF tension dominates in the Lorentz term (for $n > -1.5$) and the tension acts on the density field in the same direction as the magnetic field pressure. Yet another possibility is that the PMF tension dominates in the Lorentz term ($n > -1.5$), but the tension acts on the density field in the opposite direction from the pressure force. In these cases the PMF effects act like a gas pressure to oppose the gravitational collapse and cause the density perturbations to more slowly evolve.

On the other hand, if $s > 0$, the matter and PMF distributions could positively correlate ($s_{[\mathrm{DF}]} > 0$) and the PMF tension dominates in the Lorentz term ($n > -1.5$) while the tension acts on the density field in the opposite direction from the pressure force. Alternatively, the matter and PMF distributions could negatively correlate ($s_{[\mathrm{DF}]} < 0$) and the PMF pressure dominate in the Lorentz term ($n < -1.5$). In these cases the Lorentz force from the PMF accelerates the gravitational collapse. After decoupling, δ does not oscillate and the perturbation evolution is straightforward for all of the above cases.

V. RESULTS AND DISCUSSIONS

For completeness, in this section we briefly review the effects of the PMF on the cosmological density field fluctuations (see [37] for details). We will then illustrate that relation between σ_8 and the PMF parameters. We will show that the constraints on σ_8 from observation give a strong constraint on the PMF parameters. Since parameters of the PMF have a strong degeneracy, the existence of such a prior, can be used to effectively constrain the PMF. Also, since σ_8 is constrained by diverse observational data on linear cosmological scales, we can obtain a reliable prior for use in determining likelihood functions for the parameters of the PMF from CMB observations. The PMF effects dominate the matter power spectra for wavenumbers $k > 0.1$ Mpc^{-1}[37]. This is because the PMF energy density fluctuations depend only on the scale factor a and can survive below the Silk damping scale. Therefore, the PMF continues to source the fluctuations through the Lorentz force even below the Silk damping scale. In the case of no correlation between the PMF and the matter density fluctuations, the matter power spectrum is increased by the PMF, independently of whether the PMF pressure or tension dominates.

Here we note the different effects of the PMF on the power spectrum function $P(k)$ and the matter density fluctuation δ. While the total density fluctuation δ can be smaller or larger depending upon whether the effect of the PMF is dominated by its pressure or tension, the power spectrum function $P(k)$ always increases when the PMF does not correlate with the primordial density fluctuations. This is because $P(k) \propto \delta^2$ and is not affected by the sign of δ.

A. Effects of PMF parameters on σ_8

The alternative normalization parameter σ_8 is the root-mean-square of the matter density fluctuation in a comoving sphere of radius $8h^{-1}$ Mpc. It is given by a weighted integral of the matter power spectrum [42]. We can study the

physical processes of density field fluctuations on cosmological scales within the linear regime to determine σ_8. Recently σ_8 has been constrained by observations [58, 59, 60, 61] to be in the range $0.7 < \sigma_8 < 0.9$. From this we can obtain strong constraint for the PMF parameters by numerically calculating σ_8 under the influence of PMF effects.

We expect that the discrepancy between theoretical estimates and observational temperature fluctuations of the CMB for higher multipolarity ($\ell > 1000$) is solved by combining a PMF of strength $2.0\ nG < |B_\lambda| < 3.0\ nG$ and the SZ effects. In this case, σ_8 derived by such a field strength for the PMF is $0.77 - 0.88$. This is consistent with our assumed prior in the range σ_8 as $0.7 < \sigma_8 < 0.9$. Since σ_8 is affected by other cosmological parameters, Ω_b, $\Omega_{\rm CDM}$, $n_{\rm S}$, and $A_{\rm S}$, we should consider the degeneracy between the PMF and other cosmological parameters as mentioned above. Fortunately, these cosmological parameters are constrained by recent CMB observations on larger scales ($\ell < 1000$) [62, 63, 64], while it was shown in our previous work [21, 23, 37] that the effect of the PMF mainly affects the CMB anisotropies on smaller scales ($\ell > 1000$). Hence, we expect that the degeneracy between the PMF parameters and the other cosmological parameters is small. For this reason in the present analysis we are justified in fixing the other cosmological parameters at their best fit values.

Figure 1 shows the behavior of the PMF parameters B_λ and $n_{\rm B}$ for various constant values of σ_8 as labeled. Since the PMF power spectrum depends upon $n_{\rm B}$ from Eqs. (13-15)[25], for $n_{\rm B} > -1.5$, the PMF effects on density fluctuations for the small scales decrease with lower values for $n_{\rm B}$. While, for $n_{\rm B} < -1.5$, $P(k)_{\rm PMF} \propto B_\lambda^{[(2n_{\rm B}+3)/(n_{\rm B}+5)]}$ from Eqs. (13-15)[25]. Therefore, for a spectral index near $n_{\rm B} = -3.0$, the matter power spectrum in the presence of a PMF is smaller, and larger amplitudes of B_λ are allowed. Since the correlation effects of the PMF for negative and positive correlations change at $n_{\rm B} = -1.5$ (see Section IV), we divide the discussion below of each correlation into two parts based upon whether the spectral index is greater or less than $n_{\rm B} = -1.5$.

1. No Correlation

When there is no correlation between the PMF and the density fluctuations from primary perturbations, the third terms in both Eqs. (42) and (43) vanish and the PMF only acts to increases the total matter power spectrum. In this paper, we adopt the constraint that PMF parameters giving $\sigma_8 > 1$ are excluded by observations. Panel (a) of Figure.1 shows that a PMF amplitude of $B_\lambda \gtrsim 1$ nG is excluded when $n_{\rm B} > -0.9$. Furthermore, PMF amplitudes of $B_\lambda \gtrsim 0.11$ nG are excluded when $n_{\rm B} > 0.2$. The magnetic field strength in galaxy clusters is $\sim 1\mu$G. Therefore, if isotropic collapse is the only process which amplifies the magnetic field strength, the lower limit to the PMF is ~ 1nG at $z \sim 0$. Hence, we can obtain a strong constraint on this PMF evolution model for a PMF spectral index in the range $n_{\rm B} < -0.9$.

2. Negative Correlation

For the case of negative correlations, the pressure of the PMF dominates for $n_{\rm B} < -1.5$, and the PMF causes an increase in the density fluctuations. For $n_{\rm B} > -1.5$, however, the tension of the PMF dominates, and the PMF causes a decrease in the density fluctuations. These behaviors can be traced to the third terms in each Eqs.(42) and (43). Using the allowed range of PMF parameters as mentioned above, a PMF of $B_\lambda \gtrsim 1$ nG is excluded for $n_{\rm B} > -0.81$, and a PMF of $B_\lambda \gtrsim 0.11$ nG is excluded for $n_{\rm B} > 0.26$.

3. Positive Correlation

When there is a positive correlation, the pressure of the PMF dominates for $n_{\rm B} < -1.5$, and the PMF leads to a decrease in the density fluctuations. On the other hand, the tension of the PMF dominates for $n_{\rm B} > -1.5$. In this case the PMF causes an increase in the density fluctuations. We can attribute these behaviors again to the third terms in Eqs. (42) and (43). Using the allowed range of PMF parameters as noted above, a PMF of $B_\lambda \gtrsim 1$ nG is excluded for $n_{\rm B} > -0.94$, and a PMF of $B_\lambda \gtrsim 0.11$ nG is excluded for $n_{\rm B} > 0.13$

For both negative correlations with $n_{\rm B} > -1.5$ and positive correlations with $n_{\rm B} < -1.5$, the PMF decreases the total matter power spectrum until the strength of the PMF effect is comparable to the primary matter power spectrum. Beyond this point, the density fluctuations from the PMF exceed the density fluctuations from the primary power spectrum and the PMF effect dominates the total matter power spectrum. The amplitude of the total matter power spectra when a PMF is present is greater than the total matter power spectrum without the presence of a PMF.

In these cases, we obtain different constraints on the strength of the PMF for each value of $n_{\rm B}$ from the other models. We assume that ranges of the PMF parameters giving $\sigma_8 < 0.6$ are excluded by observations. Panel (c-1) of Figure.1 shows the excluded range of the PMF parameters within the contour defined by $\sigma_8 = 0.6$.

Our result is consistent with previous constraints on PMF parameters [22, 56], and our new more precise constraints obtained from the matter power spectrum are independent of previous methods to constrain the PMF parameters. Hence, we can now constrain more precisely the physical processes by which a PMF affects the evolution of structure on cosmological scales.

VI. SUMMARY

A primordial magnetic field (PMF) affects the evolution of density field fluctuations in the early universe. Therefore, we can constrain PMF parameters, e.g. the PMF amplitude B_λ and power spectral index $n_{\rm B}$, by comparing a theoretical calculation of the density field fluctuations affected by a PMF and observational data, e.g. a number density fluctuation of galaxies as indicated by the σ_8 parameter. We have illustrated the relation between σ_8 and the PMF parameters. We have shown that the observed range of σ_8 gives a strong limit on the PMF parameters given by $n_{\rm B} < 0.9$ (no correlation case), < -0.86 (negative correlation case), and < 0.94 (positive correlation case) for $B_\lambda > 1$nG. We have also shown how the PMF parameters which were constrained by previous methods are affected by our new constraint from σ_8. Since density field fluctuations are the origin of the LSS on cosmological scales, we can use this as a prior to study the physical processes of the PMF and to place better constraints on the PMF parameters.

Acknowledgments

We acknowledge Drs. K. Umezu, and H. Hanayama for their valuable discussions. K.I. acknowledges the support by Grants-in-Aid for JSPS Fellows. This work has been supported in part by Grants-in-Aid for Scientific Research (17540275, 20244035) of the Ministry of Education, Culture, Sports, Science and Technology of Japan, and the Mitsubishi Foundation. This work is also supported by the JSPS Core-to-Core Program, International Research Network for Exotic Femto Systems (EFES). Work at UND supported in part by the US Department of Energy under research grant DE-FG02-95-ER40934.

[1] P. P. Kronberg, J. J. Perry, and E. L. H. Zukowski, ApJ. **387**, 528 (1992).
[2] A. M. Wolfe, K. M. Lanzetta, and A. L. Oren, ApJ. **388**, 17 (1992).
[3] T. E. Clarke, P. P. Kronberg, and H. Boehringer, Astrophys. J. **547**, L111 (2001).
[4] Y. Xu, P. P. Kronberg, S. Habib, and Q. W. Dufton, Astrophys. J. **637**, 19 (2006).
[5] G. Betschart, P. K. S. Dunsby, and M. Marklund, Class. Quant. Grav. **21**, 2115 (2004).
[6] D. Boyanovsky, M. Simionato, and H. J. de Vega, Phys. Rev. **D 67**, 023502 (2003).
[7] M. Bruni, R. Maartens, and C. G. Tsagas, Mon. Not. Roy. Astron. Soc. **338**, 785 (2003).
[8] K. Bamba and J. Yokoyama, Phys. Rev. **D 70**, 083508 (2004).
[9] Z. Berezhiani and A. D. Dolgov, Astropart. Phys. **21**, 59 (2004).
[10] A. D. Dolgov (2005), astro-ph/0503447.
[11] H. Hanayama et al., Astrophys. J. **633**, 941 (2005).
[12] K. Takahashi, K. Ichiki, H. Ohno, and H. Hanayama, Phys. Rev. Lett. **95**, 121301 (2005).
[13] K. Ichiki, K. Takahashi, H. Ohno, H. Hanayama, and N. Sugiyama, Science **311**, 827 (2006).
[14] J. M. Quashnock, A. Loeb, and D. N. Spergel, Astrophys. J. Lett. **344**, L49 (1989).
[15] D. Grasso and H. R. Rubinstein, Phys. Rept. **348**, 163 (2001).
[16] K. Takahashi, K. Ichiki, and N. Sugiyama, Phys. Rev. **D77**, 124028 (2008).
[17] K. Ichiki, K. Takahashi, N. Sugiyama, H. Hanayama, and H. Ohno (2007), astro-ph/0701329.
[18] K. Takahashi (2008), 0804.3578.
[19] D. G. Yamazaki, K. Ichiki, and T. Kajino, Nuclear Physics A **758**, 791 (2005).
[20] D. G. Yamazaki, K. Ichiki, and T. Kajino, Astrophys. J. **625**, L1 (2005).
[21] D. G. Yamazaki, K. Ichiki, T. Kajino, and G. J. Mathews, Astrophys. J. **646**, 719 (2006).
[22] D. G. Yamazaki, K. Ichiki, T. Kajino, and G. J. Mathews, PoS(NIC-IX). p. 194 (2006).
[23] D. G. Yamazaki, K. Ichiki, T. Kajino, and G. J. Mathews, Phys. Rev. **D 77**, 043005 (2008).
[24] K. Kojima, K. Ichiki, D. G. Yamazaki, T. Kajino, and G. J. Mathews, Phys. Rev. **D78**, 045010 (2008).
[25] A. Mack, T. Kahniashvili, and A. Kosowsky, Phys. Rev. **D 65**, 123004 (2002).
[26] K. Subramanian and J. D. Barrow, Phys. Rev. Lett. **81**, 3575 (1998).



FIG. 1: Curves of constant values for σ_8 in the parameter plane of PMF amplitude B_λ vs. spectral index $n_{\rm B}$. Blue dashed, green dotted, red bold, black dotted thin, and azure dashed thin curves show constant values of σ_8 =1.0, 0.9, 0.8, 0.7 and 0.6, respectively.

[27] K. Subramanian and J. D. Barrow, Mon. Not. Roy. Astron. Soc. **335**, L57 (2002).
[28] A. Lewis, Phys. Rev. **D 70**, 043011 (2004).
[29] A. Kosowsky, T. Kahniashvili, G. Lavrelashvili, and B. Ratra, Phys. Rev. **D 71**, 043006 (2005).
[30] T. Kahniashvili and B. Ratra, Phys. Rev. **D 71**, 103006 (2005).
[31] T. Kahniashvili and B. Ratra, Phys. Rev. **D75**, 023002 (2007).
[32] T. Kahniashvili, G. Lavrelashvili, and B. Ratra, Phys. Rev. **D78**, 063012 (2008).
[33] A. Challinor, Lect. Notes Phys. **653**, 71 (2004).

[34] R. Gopal and S. K. Sethi, Phys. Rev. **D 72**, 103003 (2005).
[35] M. Giovannini, Phys. Rev. **D 74**, 063002 (2006).
[36] M. Giovannini and K. E. Kunze, Phys. Rev. **D78**, 023010 (2008).
[37] D. G. Yamazaki, K. Ichiki, K. I. Umezu, and H. Hanayama, Phys. Rev. **D 74**, 123518 (2006).
[38] C. G. Tsagas and R. Maartens, Phys. Rev. **D 61**, 083519 (2000).
[39] M. Giovannini, Phys. Rev. **D 70**, 123507 (2004).
[40] H. Tashiro, N. Sugiyama, and R. Banerjee, Phys. Rev. **D 73**, 023002 (2006).
[41] S. K. Sethi, B. B. Nath, and K. Subramanian (2008), arXiv:0804.3473.
[42] P. J. E. Peebles, *The Large-Scale Structure of the Universe* (Princeton University Press, 1980).
[43] J. Dunkley et al. (WMAP) (2008), arXiv:0803.0586.
[44] R. Durrer, P. G. Ferreira, and T. Kahniashvili, Phys. Rev. **D 61**, 043001 (2000).
[45] K. Subramanian and J. D. Barrow, Phys. Rev. **D 58**, 083502 (1998).
[46] R. Banerjee and K. Jedamzik, Phys. Rev. **D 70**, 123003 (2004).
[47] S. Koh and C. H. Lee, Phys. Rev. **D 62**, 083509 (2000).
[48] I. Brown and R. Crittenden, Phys. Rev. **D 72**, 063002 (2005).
[49] D. G. Yamazaki, preparing , (2008).
[50] T. Padmanabhan, *Structure formation in the universe* (Cambridge University Press, 1993).
[51] C.-P. Ma and E. Bertschinger, Astrophys. J. **455**, 7 (1995).
[52] W. Hu and M. J. White, Phys. Rev. **D 56**, 596 (1997).
[53] W. Hu, U. Seljak, M. J. White, and M. Zaldarriaga, Phys. Rev. **D 57**, 3290 (1998).
[54] S. Dodelson, *Modern Cosmology* (Academic Press, 2003).
[55] S. Chandrasekhar, *Hydrodynamic and Hydromagnetic Stability* (Claredon Press, Oxford, 1961).
[56] C. Caprini and R. Durrer, Phys. Rev. **D 65**, 023517 (2001).
[57] H. Tashiro and N. Sugiyama, Mon. Not. Roy. Astron. Soc. **368**, 965 (2006).
[58] S. Cole et al. (The 2dFGRS), Mon. Not. Roy. Astron. Soc. **362**, 505 (2005), astro-ph/0501174.
[59] M. Tegmark et al. (SDSS), Phys. Rev. **D74**, 123507 (2006), astro-ph/0608632.
[60] E. Rozo et al. (2007), astro-ph/0703571.
[61] A. J. Ross, R. J. Brunner, and A. D. Myers (2008), 0804.3325.
[62] D. N. Spergel et al. (2006), astro-ph/0603449.
[63] G. Hinshaw et al. (2006), astro-ph/0603451.
[64] L. Page et al. (2006), astro-ph/0603450.